

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

PREVEXXION RN+HVT+IBD concentrate and solvent for suspension for injection

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose (0.2 ml for subcutaneous or 0.05 ml for *in ovo*) of the vaccine suspension contains:

Active substances:

Marek's disease (MD) virus, serotype 1,
strain RN1250 (cell-associated), live:

2.9 to 3.9 log₁₀ PFU*

Turkey herpesvirus (HVT), strain vHVT013-69 (cell-associated),
expressing the VP2 protein gene of infectious bursal disease (IBD), strain Faragher
52/70 virus, live:

3.6 to 4.4 log₁₀ PFU*

*PFU: plaque-forming units

Excipients:

Qualitative composition of excipients and other constituents
Vaccine concentrate:
Dimethyl sulfoxide
199 Earle medium
Sodium hydrogen carbonate
Hydrochloric acid
Water for injections
Solvent:
Sucrose
Casein hydrolysate
Phenolsulfonphthalein (Phenol red)
Dipotassium phosphate
Potassium dihydrogen phosphate
Sodium hydroxide or hydrochloric acid
Water for injections

Concentrate: yellow to reddish pink opalescent homogeneous suspension.
Solvent: red-orange limpid solution.

3. CLINICAL INFORMATION

3.1 Target species

Chickens

3.2 Indications for use for each target species

For active immunisation of one-day-old chicks or 18-day-old embryonated chicken eggs:

- to prevent mortality and clinical signs and reduce lesions caused by MD virus (including very virulent MD virus), and
- to prevent mortality and clinical signs and lesions caused by IBD (also known as Gumboro disease) virus.

Onset of immunity: MD: 5 days post-hatch.
IBD: 14 days post-hatch (subcutaneous) or 28 days post-hatch (*in ovo*).

Duration of immunity: MD: A single vaccination is sufficient to provide protection for the entire risk period.
IBD: 10 weeks post-hatch.

3.3 Contraindications

None.

3.4 Special warnings

Vaccinate healthy animals only.

Chickens with maternally derived antibodies against MD when vaccinated with this veterinary medicinal product, may have a delayed onset of immunity against IBD.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Apply the usual aseptic precautions to all administration procedures.

As this is a live vaccine, both vaccine strains may be excreted from vaccinated birds.

The RN1250 vaccine strain has not been shown to spread in experimental conditions.

The vHVT013-69 vaccine strain may be spread to unvaccinated chickens and turkeys.

Appropriate veterinary and husbandry measures should be taken to avoid spread of the vaccine strains to unvaccinated chickens, turkeys and other susceptible species.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Personal protective equipment consisting of gloves, spectacles and boots should be worn when handling the veterinary medicinal product, before withdrawing from liquid nitrogen, during the ampoule thawing and opening operations. Frozen glass ampoules may explode during sudden temperature changes. Store and use liquid nitrogen only in a dry and well-ventilated place. Inhalation of the liquid nitrogen is dangerous.

Special precautions for the protection of the environment:
Not applicable.

3.6 Adverse events

Chickens:
None.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

This veterinary medicinal product is designed for one-day-old chicks and 18-day-old embryonated chicken eggs, therefore the safety of the veterinary medicinal product has not been established during lay.

3.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case-by-case basis.

3.9 Administration routes and dosage

Subcutaneous and *in ovo* use.

Preparation of the vaccine suspension:

- Wear protective gloves, spectacles and boots during the ampoule thawing and opening operations. The handling of liquid nitrogen should take place in a well-ventilated area.
- Preparation of the vaccine shall be planned before the ampoules are taken from the liquid nitrogen. The exact amount of vaccine ampoules and amount of solvent needed shall be calculated first according to the table below provided as example:

Solvent bag	Number of vaccine ampoules (subcutaneous use)	Number of vaccine ampoules (<i>in ovo</i> use)
1 bag of 200 ml solvent	1 ampoule (1 000 doses)	4 ampoules (1 000 doses) or 2 ampoules (2 000 doses) or 1 ampoule (4 000 doses)
1 bag of 400 ml solvent	2 ampoules (1 000 doses) or 1 ampoule (2 000 doses)	8 ampoules (1 000 doses) or 4 ampoules (2 000 doses) or 2 ampoules (4 000 doses)
1 bag of 800 ml solvent	4 ampoules (1 000 doses) or 2 ampoules (2 000 doses) or 1 ampoule (4 000 doses)	16 ampoules (1 000 doses) or 8 ampoules (2 000 doses) or 4 ampoules (4 000 doses)

- Remove from the liquid nitrogen container only those ampoules, which are to be used immediately.
- Thaw the contents of the ampoules rapidly by gentle agitation in water at 25 °C – 30 °C. The thawing process should not exceed 90 seconds. Proceed immediately to the next step.
- As soon as they are thawed, wipe the ampoules with a clean paper towel and then open them while holding them at arm's length (in order to prevent injury if any ampoule breaks).
- Select an appropriately sized sterile syringe to withdraw the vaccine from all the ampoules that are thawed, and fit it with a needle of 18 gauge or larger.
- Tear the overpouch on the solvent bag, and then gently insert the syringe needle through the septum of one of the bag-connecting tubes and withdraw 2 ml of solvent.
- Then draw up the complete contents of all the thawed ampoules into the syringe. Do this by slowly drawing up the contents from each ampoule by gently tilting the ampoule forward and inserting the needle with the bevel edge facing downwards towards the bottom of the ampoule. Continue until all the vaccine is drawn out of the ampoule.
- Transfer the syringe contents into the solvent bag (do not use the solvent if it is cloudy).
- Gently mix the vaccine in the solvent bag by moving the bag back and forth.
- It is important to rinse the ampoules and ampoule tips. To do this, draw up a small volume of the solvent containing the vaccine into the syringe. Then slowly fill the ampoule bodies and tips with it. Withdraw the content from the ampoule bodies and tips, and inject it back into the solvent bag.
- Repeat this rinsing operation once.

- Repeat the thawing, opening, transfer and rinsing operations for the appropriate number of ampoules to be diluted in the solvent bag.
- The vaccine is ready for use and should be mixed by gentle agitation and used immediately. During vaccination, gently swirl the bag frequently to ensure the vaccine remains homogenously mixed.
- The vaccine is a clear, red-orange coloured suspension for injection to be used within two hours. Do not freeze it under any circumstances. Do not re-use opened containers of vaccine.

Posology:

One single injection of 0.2 ml per one-day-old chick or 0.05 ml per 18-day-old embryonated chicken egg.

Method of administration:

The vaccine must be administered by subcutaneous injection in the neck or by *in ovo* injection.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

A limited and transient effect on growth was observed when 10-fold maximum release dose was administered subcutaneously to White Leghorn specified pathogen free chickens.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Zero days.

4. IMMUNOLOGICAL INFORMATION

4.1 ATCvet code: QI01AD15

The vaccine contains the recombinant viruses RN1250 and vHVT013-69 within chicken embryo cells.

The RN1250 virus is an engineered MD virus composed of three serotype 1 strains. Its genome also contains long terminal repeats of reticuloendotheliosis virus.

The vHVT013-69 virus is a recombinant HVT expressing the protective antigen (VP2) of the IBD virus strain Faragher 52/70.

The vaccine induces an active immunity and a serological response against Marek's disease and IBD in chickens.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

Shelf life of the solvent as packaged for sale: 3 years.

Shelf life after dilution according to directions: 2 hours at a temperature below 25 °C.

5.3 Special precautions for storage

Vaccine concentrate:

Store and transport frozen in liquid nitrogen.

The liquid nitrogen containers must be checked regularly for liquid nitrogen level and must be refilled as needed.

Discard any ampoules that have been accidentally thawed.

Solvent:

Store below 30 °C. Do not freeze. Protect from light.

5.4 Nature and composition of immediate packaging

Vaccine concentrate:

- Type I glass ampoule of 1 000 doses of vaccine.
- Type I glass ampoule of 2 000 doses of vaccine.
- Type I glass ampoule of 4 000 doses of vaccine.

Each ampoule is placed on carriers which are stored in canisters. The canisters are further stored in liquid nitrogen containers.

Solvent:

- Polyvinylchloride bag containing 200 ml, 400 ml, 600 ml, 800 ml, 1 000 ml, 1 200 ml, 1 600 ml, 1 800 ml or 2 400 ml.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Boehringer Ingelheim Vetmedica GmbH

7. MARKETING AUTHORISATION NUMBER

Vm 61700/5041

8. DATE OF FIRST AUTHORISATION

20 July 2020

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

September 2025

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

Gavin Hall
Approved: 22 December 2025